

Impact of biochar on crop yields in temperate soils

A summary of field trial data looking at application rates of biochar in temperate regions - and subsequent positive impacts on crop yield.

Type of crop	Location	Type of soils	Authors	Quantity of biochar (ton/ha)	Yield increases %
Canola	Saskatchewan, Canada	Brown loam	Pervej-Ahmed et al.	1	20%
Wheat	Postoia, Italy	Silty loam	Vaccari et al.	30	33%
Rice	Northern Italy	Aguic hapludalf	Lugato et al.	40	36%
Rice	Shenyang, China	Sandy loam	Zhang et al.	30	40%
Radish	NSW, Australia	Shromosol	Chan et al.	10	42%
Soyabean	Gifu, Japan	N/A	Tagoe et al.	3.8	43%
Barley	United Kingdom	Light soil	Gathorne-Hardy et al.	20	43%
Quinoa	Germany	Sandy loam brown earth	Kamman et al.	100	44%
Citrus	Matsuyama, Japan	N/A	Ishii et al.	83.5	57%
Maize	Yingtian, China	Ultisol	Peng et al.	2.4	64%
Sweet pepper	Israel	Commercial soilless mixture	Graber et al.	1%	79%
Choux	Nanjing, China	Fimi-orthic anthrosol	Jia et al.	30	96%
Maize	Tottori, Japan	Sandy soil	Uzoma et al.	15	150%
Wheat	NSW, Australia	Ferralsol	Van Zwieten	2%	170%

For more information please visit www.carbongold.com

Impact of biochar on crop yields in tropical soils

A summary of field trial data looking at application rates of biochar in tropical regions - and subsequent positive impacts on crop yield.

Type of crop	Location	Type of soils	Authors	Quantity of biochar (ton/ha)	Yield increases %
Casava	Malang, Indonesia	Clay loam	Islami et al.	15	32%
Onion	Senegal	N/A	Pro-Natura	10	50%
Peanuts	Malang, Indonesia	Clay loam	Islami et al.	15	54%
Rice	Empretring, Indonesia	N/A	Zaitun et al.	10	57%
Rice	Houay-Khot, Nord du Laos	Upland	Asai et al.	8	70%
Maize	Llanos Orientales, Colombia	Savanna oxisol	Major et al.	8	71%
Maize	Vihiga, western Kenya	Highly degraded ultisol	Kimetu et al.	6	71%
Rice	Manuas, Brazil	Xanthic ferralsol / laterite	Steiner et al.	11	73%
Sugarcane	Okinawa, Japan	Shimajiri maji (clay)	Chen et al.	7.2	78%
Rice	Sungai Kakap, Indonesia	Acid sulphate soil	Masulili et al.	10	93%
Cotton	Midjil Mandal, Andhra Pradesh, India	Alkaline	Reddy	3.75	100%
Maize	Llanos Orientales, Colombia	Savanna oxisol	Major et al.	20	140%
Cowpea	Gifu, Japan	Sandy loam	Tagoe et al.	N/A	146%
Tomato	Kade, Ghana	Forest ochrosol	Effah et al.	7	177%
Cabbage	Siam Reap, Cambodia	Sandy acidic	Carter et al.	100	750%

For more information please visit www.carbongold.com